

CLENBUTEROL HYDROCHLORIDE

DESCRIPTION

Clenbuterol is a prescribed Asthma medication which is catabolic to fat and anabolic to muscle. It is a Beta-2 adrenergic agonist. This means it acts just like adrenaline (epinephrine) at a certain type of receptor named Beta-2.

Clenbuterol belongs to a broad group of drugs known as sympathomimetics. These drugs affect that sympathetic nervous system in a wide number of ways, largely mediated by the distribution of adrenoceptors. There are actually nine different types of these receptors in the body, which are classified as either alpha or beta and further subcategorized by type number. Depending on the specific affinities of these agents for the various receptors, they can potentially be used in the treatment of conditions such as asthma, hypertension, cardiovascular shock, arrhythmias, migraine headaches and anaphylactic shock.

Chemical: Clenbuterol Hydrochloride
Additional Names: 4-Amino-alpha-[(tert-butylamino)methyl]-3,5-dichlorobenzyl alcohol monohydrochloride
Molecular Formula: C₁₂H₁₈Cl₂N₂O.HCl
Molecular Weight: 313.65

Prescription Medicine

CONTRAINDICATIONS

Individuals who have been diagnosed of cardio-vascular problems should not take Clenbuterol.

DOSAGE AND USAGE

Clenbuterol Hydrochloride tablets should be orally consumed.
Adult dose: 60-120 mcgs per day, divided throughout the day.

SIDE EFFECTS AND SPECIAL PRECAUTIONS

Nervousness, tremors of the hands, headaches, early onset asthma and insomnia.

As long as the electrolyte status is maintained (esp. potassium) - clenbuterol does not cause too many problems. Muscle cramps are the most common side effect. If you have a job that requires lots of manual dexterity, then the usage of the clenbuterol can be highly problematic. The tremor it gives in the hands can make steady fine movements virtually impossible. Fortunately, this effect often diminishes over a week or two of continued usage. Clenbuterol can have some serious side effects (like sudden death) at very high dosages. Blood pressure should be monitored constantly when using Clenbuterol. The dosage should be based on individuals blood pressure and resting pulse.

STORAGE

Store in a cool dry place (30 C ± 2 C). Protect from light.

PRESENTATION:

0.04mg tablets in blister packs of 10 tablets – 10 blisters per box (100 tablets).

AURORA REMEDIES, SINGAPORE

www.aurora-remedies.com